



SURGICAL SERVICES - NORTH AMERICA

Olfactory Bulbectomy

Surgery Code: OLFACTOREX

The olfactory bulbectomy is of benefit to preclinical or research studies involving the impaired sense of smell.

Animal Models

Typical selections are listed below; however, choices for strain, age and weight may be limited due to model anatomy and/or physiological conditions.

Rats: male/female, any strain, weight ≥ 200 g

Procedure Details

- Perioperative care: Please view our Pre- and Postoperative Care Sheet, which can be found at www.criver.com/opcare.
- · Housing: The animals can be group housed.
- Diet: No special diet is required.
- Postoperative holding period: At a minimum, post-op animals are held 2 days, with the majority of animals shipping within 7 days of surgery.
- Maintenance: Wound clips should be removed 7-10 days after surgery.

Surgical Summary

The animal is placed in the stereotaxic apparatus and skin incision is made on the skull. A hole is drilled into the skull to access the olfactory bulbs. The bases of the bulbs are separated and the olfactory bulbs are aspirated. The skin incision is closed with wound clips or per request.

IACUC

The Charles River Institutional Animal Care and Use Committee (IACUC) governs the entire surgical process, including all anesthesia, analgesia, animal preparation and any postoperative holding in Charles River facilities prior to shipment. Review of experimental protocols, authorization to order animals that are surgically modified from Charles River, and all aspects concerning the use of animals after they arrive at the institution are the responsibility of the receiving institution's IACUC.

Contact Us

For more information, visit www.criver.com/surgery. For specific surgery-related questions, please contact our technical experts at 1.877.CRIVER.1 (1.877.274.8371) or askcharlesriver@crl.com. To place an order or get a quote, contact our Customer Service Department at 1.800.LABRATS (1.800.522.7287).